



Sonoran Desert Rapid Ecoregional Assessment Report




Final Report Organization




❑ Main Report



❑ Inserts



❑ Appendices



❑ Electronic Hosting

Inserts

Species Conservation Element

Change Agent

Agassiz's Desert Tortoise (*Gopherus agassizii*) and Morafka's Desert Tortoise (*Gopherus morafkai*)

The desert tortoise was selected as a core conservation element for the Sonoran Desert REA because it is an iconic species of the region that reflects inter-regional variability in climate, landform, and vegetation. The tortoise is a good indicator of desert condition because the two species are widely distributed across the ecoregion and, at the same time, sensitive and vulnerable to multiple disturbance factors. REA results produced maps for current status and future condition of the two populations (now species) of desert tortoise.

The desert tortoise inhabits desert environments in the Mojave and Sonoran deserts in southern California, southern Nevada, Arizona, southwestern Utah, and northwestern Mexico. Once recognized as a single species (*Gopherus agassizii*) with two recognized populations, it has recently been split into two species. The former Mojave population (Agassiz's desert tortoise) retains the name *Gopherus agassizii*; it occurs north and west of the Colorado River. It was listed as threatened in 1990–22 years after listing, the species is still declining, particularly in the western portion of its range in California (Brussard et al. 1994, Doak et al. 1994, Tracy et al. 2004). The former Sonoran population has been named Morafka's desert tortoise (*Gopherus morafkai*), distinguished from *agassizii* by its different habitat, life history traits, and DNA evidence (USGS 2011). Morafka's desert tortoise occurs east and south of the Colorado River, from Arizona into Mexico.

Agassiz's Desert Tortoise (*Gopherus agassizii*)

Agassiz's desert tortoise occurs mainly in creosote bush (*Larrea tridentata*) flats, but it is also found on sloping terrain on alluvial fans or foothills. It forages mostly on annual plants produced by winter rains. The yearly life cycle of the Agassiz's desert tortoise is heavily influenced by the annual precipitation pattern in the western Sonoran (and Mojave) Desert—precipitation that mainly falls in the winter and early spring with little or no summer precipitation (Van Devender 2006, Dickinson et al. 2006). As a result, most Agassiz's tortoise activity takes place in the spring when winter annuals and spring grasses are readily available (Nagy and Medina 1986, Brussard et al. 1994). The species faces the prospect of annual summer drought; in the hot summer months and throughout the winter, the tortoises spend many months of inactivity in burrows in estivation or hibernation without eating or drinking. In years of low winter rainfall, Agassiz's may feed on introduced annual grasses in the absence or scarcity of winter annuals (Esque 1994), and the species' summer hatchlings may not eat fresh forage until the following spring. While it is known that a diet of invasive grasses will keep tortoises alive, it is unknown if over time such a diet will keep them fit (Esque et al. 2006). Because they are at the northern limit of their range and because of their dietary restraints and restricted access to water, Agassiz's tortoise may be more vulnerable to mortality from drought, loss of condition, and other stressors than Morafka's tortoise (Peterson 1996, Offedal 2006).



Photo: Agassiz's desert tortoise, formerly Mojave desert tortoise. K. Nusser, U.S. Geological Survey

Case Study No. 1

Tamarisk (*Tamarix* spp.)

The history of the expansion of tamarisk throughout the riparian areas of the southwestern U.S. parallels the development and allocation of water resources in arid and semi-arid ecosystems in the 20th and 21st centuries. Tamarisk (or saltcedar) is an invasive shrub that has been designated as a change agent in the Sonoran Desert REA because it negatively affects aquatic resources, native riparian ecosystems, and aquatic sites of conservation concern. The name tamarisk refers to a number of related species in the genus *Tamarix* (e.g., *T. canescens*, *T. chinensis*, and *T. aphylla*) that are similar in appearance and that hybridize freely (Gaskin and Shafroth 2005). Introduced into North America at the time of Spanish settlement, the species did not become widely distributed until the 1800s. It is presently found throughout nearly all western and southwestern states (Lovich 2000). In a survey of 475 gaging stations across the western U.S., Friedman et al. (2005) found tamarisk to be the third most frequently-occurring riparian woody plant in the region.




Photo credit: Columbia University Invasive Species Summary Project



Figure 1. Current distribution of tamarisk (in blue) near the confluence of the Colorado and Gila Rivers as mapped for the Sonoran Desert REA. See Appendix XX for modeling approach and region-wide results.

Appendices



Management Questions

A

SONORAN DESERT

A. SOILS, BIOLOGICAL CRUSTS, AND FORAGE MANAGEMENT

Seq

- MQ A1. Where are soils susceptible to wind and water erosion?
- MQ A2. Where are sensitive soils (including saline, sodic, gypsiferous, shallow, low water holding capacity)?
- MQ A3. Which HMAs and allotments may experience significant effects from change agents including climate change?

B. SURFACE AND GROUNDWATER MANAGEMENT QUESTIONS

- MQ B1. Where are lotic and lentic surface waterbodies and livestock and wildlife watering tanks and artificial water bodies?
- MQ B2. Where are perennial streams and stream reaches?
- MQ B3. Where are the alluvial aquifers and their recharge areas (if known)?
- MQ B4. Where are the aquatic systems listed on 303(d) with degraded water quality or low macroinvertebrate diversity?
- MQ B5. Where are surface water flows likely to increase or decrease in the near-term, 2025 (development), and long-term, 2060 (climate change)?
- MQ B6. What is the location/distribution of these aquatic biodiversity sites?
- MQ B7. What are seasonal discharge maxima and minima for the Colorado River and major tributaries at gaging stations?

C. ECOLOGICAL SYSTEMS MANAGEMENT QUESTIONS

- MQ C1. Where are existing vegetative communities? - Sonoran Paloverde-Mixed Cacti Desert Scrub (Saguaro)
- MQ C2. Where are vegetative communities vulnerable to change agents in the future? - Sonoran Paloverde-Mixed Cacti Desert Scrub (Saguaro)
- MQ C3. What change agents have affected existing vegetation communities? - Sonoran Paloverde-Mixed Cacti Desert Scrub (Saguaro) - Historic Change
- MQ C3. What change agents have affected existing vegetation communities? - Sonoran Paloverde-Mixed Cacti Desert Scrub (Saguaro) - Recent Disturbances
- MQ C1. Where are existing vegetative communities? - Sonoran-Mojave Creosotebush-White Bursage Desert Scrub (Creosotebush)
- MQ C2. Where are vegetative communities vulnerable to change agents in the future? - Sonoran-Mojave Creosotebush-White Bursage Desert Scrub (Creosotebush)
- MQ C3. What change agents have affected existing vegetation communities? - Sonoran-Mojave Creosotebush-White Bursage Desert Scrub (Creosotebush) - Historic Change
- MQ C3. What change agents have affected existing vegetation communities? - Sonoran-Mojave Creosotebush-White Bursage Desert Scrub (Creosotebush) - Recent Disturbances
- MQ C1. Where are existing vegetative communities? - Sonoran Mid-Elevation Desert Scrub
- MQ C2. Where are vegetative communities vulnerable to change agents in the future? - Sonoran Mid-Elevation Desert Scrub
- MQ C3. What change agents have affected existing vegetation communities? - Sonoran Mid-Elevation Desert Scrub - Historic Change
- MQ C3. What change agents have affected existing vegetation communities? - Sonoran Mid-Elevation Desert Scrub - Recent Disturbances
- MQ C1. Where are existing vegetative communities? - Mogollon Chaparral
- MQ C2. Where are vegetative communities vulnerable to change agents in the future? - Mogollon Chaparral
- MQ C3. What change agents have affected existing vegetation communities? - Mogollon Chaparral - Historic Change
- MQ C3. What change agents have affected existing vegetation communities? - Mogollon Chaparral - Recent Disturbances
- MQ C1. Where are existing vegetative communities? - Apachean-Chihuahuan Mesquite Upland Scrub
- MQ C2. Where are vegetative communities vulnerable to change agents in the future? - Apachean-Chihuahuan Mesquite Upland Scrub
- MQ C3. What change agents have affected existing vegetation communities? - Apachean-Chihuahuan Mesquite Upland Scrub - Historic Change
- MQ C3. What change agents have affected existing vegetation communities? - Apachean-Chihuahuan Mesquite Upland Scrub - Recent Disturbances

D. SPECIES CONSERVATION ELEMENT MANAGEMENT QUESTIONS

- MQ D5. What is the location/distribution of terrestrial biodiversity sites? High Biodiversity Sites
- MQ D5. What is the location/distribution of terrestrial biodiversity sites? High Ecological/Cultural Sites
- MQ D5. What is the location/distribution of terrestrial biodiversity sites? National Trails
- MQ D7. Where are HMAs located?

Mammals

- MQ D1. **Mountain Lion** - What is the most current distribution of available occupied habitat (and historic occupied habitat if available), seasonal and breeding habitat, and movement corridors (as applicable)?

Management Questions

Soils and forage management questions

MQA1. Where are soils susceptible to wind and water erosion?

Methods ♦ Process Model ♦ Conceptual Model ♦ Results

MQA2. Where are sensitive soils (including saline, sodic, gypsiferous, shallow, low water holding capacity)?

Methods ♦ Process Model ♦ Conceptual Model ♦ Results

MQA3. Which HMAs and allotments may experience significant effects from change agents including climate change?

Methods ♦ Process Model ♦ Conceptual Model ♦ Results

Management Questions

Species management questions

MQD1. What is the most current distribution and status of available occupied habitat?

MQD7. What sites and movement corridors are vulnerable to change in the near term (2015) and long-term (2060) horizon?




MQJ3. Where are areas of species conservation elements potential distribution change from climate change by 2060?

Mountain Lion
Desert Bighorn Sheep
Mule Deer


Bell's Vireo
Golden Eagle
Le Conte's Thrasher
Lucy's Warbler
SW Willow Flycatcher

Lowland Leopard Frog
Desert Tortoise (agassizzi)
Desert Tortoise (morafkai)

Conservation Element



Conceptual Model



Mountain Lion
Puma concolor

Concept Model



Attributes & Indicators

Process Model



SOD – Mountain Lion Process Model

MQ D1: Mountain Lion Current Distribution and Status

All files clipped with boundary file called Outline of 5th-level HUC (10-digit) watersheds, Sonoran Desert ecoregion


Distribution

Status



Attributes and Indicators – Mountain Lion

Element	Attribute	Indicator	Poor	Fair	Good	V. Good	Citation
Mountain Lion	Prey	Ungulate density	Low	Medium	High	Very high	Julander and Jeffrey (1964)
	Habitat degradation	Road density	0.6 km/sq km	0.4	0.2	0.0	Van Dyke et al. (1986)
	Habitat	Cover & terrain	Very dense or open cover			Rugged terrain with mixed cover	Riley (1998)
	Habitat degradation	Human development	High	Moderate	Low	No	Van Dyke et al. (1986)



Electronic Hosting

DATA  BASIN

HOME | ABOUT | BLOG | HELP | DONATE | LANGUAGE: English | LOG OUT

Explore • Create • Share • Learn



Search This Group 

My Workspace

Datasets

Maps

Galleries

People

Groups

Admin

Welcome Conservation Biology Institute

 **Sonoran Desert - Rapid Ecological Assessment**

Group Home

Profile

Bookmarks

Discussions

Members (30)

Membership Requests (0)

Supporting Documents


DATA BASIN | GROUPS | SONORAN DESERT - RAPID ECOLOGICAL ... | HOME

Manage Group 

 **SONORAN DESERT - RAPID ECOLOGICAL ASSESSMENT**

Welcome. 

This group has been created to support review of input datasets, models, and results for the BLM Rapid Ecological Assessment (REA) for the Sonoran Desert ecoregion.



[Upload Home Image](#) | [Delete Home Image](#)

 Conservation Biology Institute

Messages

 **New Notifications (18)**

Create a Map

Create a Dataset

Create a Group

Create a Gallery

HOME | ABOUT | BLOG | HELP | TERMS OF USE | CONTACT US

Report a Problem 

Sonoran Desert - Rapid Ecological Assessment

Group Home

Profile

Bookmarks

Discussions

Members (30)

Membership Requests (0)

Supporting Documents



Conservation Biology Institute

Messages

New Notifications (18)

Create a Map

Import a Dataset

Create a Group

Create a Gallery

DATA BASIN | GROUPS | SONORAN DESERT - RAPID ECOLOGICAL ... | BOOKMARKS

Export List

Manage Group

Group Bookmarks

Right click folder for menu

- Group Folders
 - + Ancillary Datasets
 - + Aquatic CEs
 - + Climate Change
 - + Development
 - + Ecological Integrity
 - + Fire
 - + Invasives
 - + Species CEs
 - Terrestrial CEs
 - + Biodiversity Sites
 - CE - Soils
 - + Additional Soils
 - + MQA1 - Wind and Water Erosion
 - + MQA2 - Sensitive Soils
 - + MQA3 - Change to HMAs and A
 - + Ecological Systems
 - + Ecological/Cultural Sites
 - + NatureServe
 - + Resource Use


Sort By:
Creation Date (newest to oldest)

Items per page:
10

Add Items

Showing 1 - 10 of 20 items

SOD MQA2 - Soil Sensitivity [private]



Credits: Conservation Biology Institute
Results for soil sensitivity for the Sonoran Desert ecoregion.

13 datasets 1 comment

Created Jul 8, 2011 by
Conservation Biology Institute

view details
open live map
[+] more

HUC Results: Highly sensitive soils - Sonoran Desert ecoregion USA [private]




Credits: Wendy Peterman, Conservation Biology Institute
This dataset shows the percent of each 5th-level HUC (10-digit) watershed with highly sensitive soils. This means one or more of the following is true: the NRCS wind erodibility group is 1 or 2, Available Water Capacity is <0.05 in the top ...

Layer Package

(Page 1 of 2)


First << Previous 1 | 2 Next >> Last

Methods




- ❑ Conceptual Models
- ❑ Process Models
- ❑ Logic Models
- ❑ FRAGSTATS
- ❑ Connectivity Modeling
- ❑ Climate Modeling


Conceptual Models




Process Models



Logic Models




Fuzzy Logic




Thresholds for Terrestrial Landscape Intactness

Item	Data Type	Data Range	True Threshold	False Threshold
Fire Regime	Percent Area	0-100	7	100
Invasive Grasses & Tamarisk	Percent Area	0-88	0	33
Linear Development	Density	0-18	0	2.5
Urban Percent	Percent Area	0-99	0	15
Agriculture Percent	Percent Area	0-90	0	20
Energy & Mining Development	Number	0-37	0	1.25
Number of Patches	Number	1-1,455	0	850
Mean Nearest Neighbor	Distance	60-272	59	180
Percent Natural Core Area	Percent Area	.56-95	100	20

Thresholds for Terrestrial Landscape Intactness



Thresholds for Terrestrial Landscape Intactness










Terrestrial Landscape Intactness Value Ranges

Intactness Value	Legend
-1.000 to -0.750	Very Low
-0.750 to -0.500	Low
-0.500 to 0.000	Moderately Low
0.000 to 0.500	Moderately High
0.500 to 0.750	High
0.750 to 1.000	Very High


Habitat Fragmentation - FRAGSTATS

4km and HUC







Fire Modeling










Climate Change Modeling



Climate Change Modeling







Distribution and Status



Species CEs	Total Distribution Area	Percent of Ecoregion
Bell's Vireo (<i>Vireo bellii</i> and <i>Vireo bellii pusillus</i>)	2,821,371	8.08%
Desert Bighorn Sheep (<i>Ovis canadensis nelsoni</i>)	7,862,801	22.52%
Desert tortoise (<i>Gopherus agassizii</i>)	3,180,566	9.11%
Desert tortoise (<i>Gopherus morafkai</i>)	6,950,996	19.91%
Golden Eagle (<i>Aquila chrysaetos</i>)	17,257,325	49.42%
Le Conte's Thrasher (<i>Toxostoma lecontei</i>)	9,771,582	27.99%
Lowland Leopard Frog (<i>Lithobates yavapaiensis</i>)	678,014	1.2%
Lucy's Warbler (<i>Oreothlypis luciae</i>)	13,753,047	39.39%
Mountain Lion (<i>Puma concolor</i>)	13,893,442	39.79%
Mule Deer (<i>Odocoileus hemionus</i>)	26,845,687	76.89%
SW Willow Flycatcher (<i>Empidonax traillii extimus</i>)	139,328	0.40%



Current Terrestrial Intactness



Current Aquatic Intactness

















Mountain Lion





Mountain Lion





Mule Deer





Desert Bighorn Sheep





Desert Tortoise (*agassizii*)





Desert Tortoise (*morafkai*)





Lowland Leopard Frog





Lucy's Warbler





Golden Eagle





LeConte's Thrasher



Bell's Vireo




Southwest Willow Flycatcher




Distribution and Status Vegetation Communities

Sonora-Mojave Creosotebush-White Bursage Desert Scrub



Sonoran Paloverde-Mixed Cacti Desert Scrub



Vegetation Community	NatureServe Only	LANDFIRE Only	Both	Percent Overlap
Sonora-Mojave Creosotebush-White Bursage Desert Scrub	5,361,364	1,417,474	4,823,357	41.6
Sonoran Paloverde-Mixed Cacti Desert Scrub	1,796,793	5,332,340	7,373,363	50.84
Riparian Vegetation	1,599,887			

Status of Vegetation Communities

☐ Historic Conversion



☐ Recent Disturbance




☐ Current Setting




EVT versus BsP

Sonora-Mojave Creosotebush-White Bursage Desert Scrub




Sonoran Paloverde-Mixed Cacti Desert Scrub




Historic Disturbance

Sonora-Mojave Creosotebush-White Bursage Desert Scrub




Sonoran Paloverde-Mixed Cacti Desert Scrub




Recent Disturbance

Sonora-Mojave Creosotebush-White Bursage Desert Scrub



Sonoran Paloverde-Mixed Cacti Desert Scrub



Historic Disturbance



Vegetation Community	Total BpS Area	Urban & Roads	Agriculture	Invasives	Unchar Native Veg	Total Changed	Percent
Sonora-Mojave Creosotebush-White Bursage Desert Scrub	7,857,699	429,263	432,831	2,909,387	273,834	4,045,315	51.48%
Sonoran Paloverde-Mixed Cacti Desert Scrub	15,730,037	1,255,201	672,008	2,345,345	429,066	4,701,621	29.89%
Totals	23,587,737	1,684,464	1,104,839	5,254,733	702,900	8,746,936	37.08%

Recent Disturbance



Vegetation Community	Total BpS Area	Fire	Mechanical	Other	Total Disturbed	Percent
Sonora-Mojave Creosotebush-White Bursage Desert Scrub	7,857,699	85,434	0	80	85,514	1.09%
Sonoran Paloverde-Mixed Cacti Desert Scrub	15,730,037	212,197	0	18	212,216	1.35%
Totals	23,587,737	297,631	0	98	297,730	1.26%

Current Setting

Sonora-Mojave Creosotebush-White Bursage Desert Scrub - LANDFIRE





Sonora-Mojave Creosotebush-White Bursage Desert Scrub - NatureServe





Current Setting



Sonoran Paloverde-Mixed Cacti Desert Scrub - LANDFIRE




Sonoran Paloverde-Mixed Cacti Desert Scrub - NatureServe





Riparian Vegetation



Designated Lands
















Designated Lands




Designated Lands

Designation Category	Very High	High	Moderately High	Moderately Low	Low	Very Low	Total Area (acres)
Wilderness Area	1,812,561	1,632,927	1,011,143	451,464	55,862	21,933	4,985,891
Wilderness Study Area	0	0	6,620	55,984	2,525	2,272	67,401
National Park	19,819	83,040	42,873	14,441	7,178	6,979	174,330
National Monument	127,785	148,652	158,627	68,922	24,265	6,305	534,556
National Conservation Area	0	6,779	21,819	7,638	95	0	36,331
National Wildlife Refuge	91,295	95,602	51,145	61,499	39,637	31,834	371,011
Area of Critical Conservation Concern	257,951	409,293	503,740	174,291	95,838	41,669	1,482,783
State Park	1,151	35,434	152,845	212,951	126,418	105,968	634,768
State Wildlife Management Area	1,202	288	4,250	3,126	4	4,819	13,689
Roadless Area	11,564	32,957	65,350	5,908	2,526	0	118,306
Other Protected Lands	71,122	153,511	177,174	243,082	43,660	97,114	785,663
Totals	2,394,451	2,598,484	2,195,588	1,299,307	398,008	318,893	9,204,730
Designation Category	Very High	High	Moderately High	Moderately Low	Low	Very Low	Total Area (miles)
National Historic and Scenic Trail	28	97	179	160	69	282	815
Wild and Scenic River	0	1	6	0	2	0	9
Totals	28	99	185	160	71	282	823


Designated Lands




Terrestrial Connectivity




Change Agent - Current Invasives




Change Agent - Current Invasives




Change Agent – Fire




Change Agent – Fire







Change Agents - Fire



Change Agent – Current Development



Change Agent – Current Development



Change Agent – Current Development

